arXiv:1612.05931v2 [math.NT] 19 Jan 2017

EXISTENCE RESULTS ON $K-$NORMAL ELEMENTS OVER FINITE FIELDS

LUCAS REIS

Abstract. An element $\alpha \in \mathbb{F}_{q^n}$ is said to be normal over $\mathbb{F}_q$ if the set $\{\alpha, \alpha^q, \cdots \alpha^{q^{n-1}}\}$ is a basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. As a generalization of normal elements, Huczynska, Mullen, Panario and Thomson (2013) introduce and characterize $k-$normal elements, proposing many problems based on theoretical considerations and computational tests. Motivated by this work, in this paper we discuss two of those problems, concerning the existence of high-order and primitive $k-$normal elements in finite field extensions.

1. Introduction

Let q be a prime power, n a positive integer and $\mathbb{F}_{q^n}$ the finite field with q^n elements. An element $\alpha \in \mathbb{F}_{q^n}$ is said to be normal over $\mathbb{F}_q$ if the set $A = \{\alpha, \alpha^q, \cdots, \alpha^{q^{n-1}}\}$ is a basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$; A is called a normal basis. Due to their efficiency in exponentiations in finite fields, normal bases are frequently used in cryptography and computer algebra systems; $q-$th powers in $\mathbb{F}_{q^n}$ correspond to a cyclic shift in the elements of a normal basis. The existence of normal basis over any finite field extension is known, as well as the existence of normal basis composed by *primitive elements*, i.e., generators of the multiplicative group $\mathbb{F}_{q^n}^*$. For more details, see [8].

It is well known ([9], Theorem 2.39) that $\alpha \in \mathbb{F}_{q^n}$ is normal over $\mathbb{F}_q$ if and only if the polynomials $g_\alpha(x) := \sum_{i=0}^{n-1} \alpha^{q^i} \cdot x^{n-1-i}$ and $x^n - 1$ are relatively prime over $\mathbb{F}_{q^n}$. Motivated by this fact, in [7], the authors introduce $k-$normal elements: for $0 \leq k \leq n-1$, an element $\alpha \in \mathbb{F}_{q^n}^*$ is said to be $k-$normal over $\mathbb{F}_q$ if the greatest common divisor of $g_\alpha(x)$ and $x^n - 1$ over $\mathbb{F}_{q^n}$ has degree k. In this context, $0-$normal elements correspond to normal elements in the usual sense. In [7] the authors give a characterization of $k-$normal elements and some bounds for their number. Also, they discuss the existence of primitive $1-$normal elements and propose some problems arising from theoretical and practical considerations:

Date: January 20, 2017.

2010 *Mathematics Subject Classification.* 12E20 (primary), 11T30 and 11T06 (secondary).

Key words and phrases. Normal basis, $k-$normal elements, Primitive elements, Elements of high order.

- Determine the pairs (n,k) such that there exist primitive $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$,
- Determine the existence of high-order $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.

In this paper we discuss the previous two problems. The paper is structured as follows. In Section 2 we present some definitions and useful preliminary results. In Section 3 we obtain an existence result for primitive $k-$normal elements, extending the previous work in [7]. Finally, in Section 4 we prove that if there exist $k-$normal elements of $\mathbb{F}_{q^n}$, then at least one of them has multiplicative order bounded below by $q^{n(1-\varepsilon(n,q))-k}$, where $\varepsilon(n,q)\to 0$ as q goes to infinity. Also, in Section 4, we give sufficient conditions for the existence of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ for any $1\leq k\leq n-1$.

2. Preliminaries

The following results and definitions concerning $k-$normal elements are contained in [7] and the further results are quite standard. For this reason we just summarize them.

2.1. k-normal elements.

Definition 2.1. *Let $\alpha\in\mathbb{F}_{q^n}^*$ and $g_\alpha(x)=\sum_{i=0}^{n-1}\alpha^{q^i}\cdot x^{n-1-i}$. We say that α is $k-$normal over $\mathbb{F}_q$ if the greatest common divisor of x^n-1 and $g_\alpha(x)$ over $\mathbb{F}_{q^n}$ has degree k.*

Definition 2.2. *The $\mathbb{F}_q-$Order of $\alpha\in\mathbb{F}_{q^n}$ is the (unique) monic polynomial $f(x)=\sum_{i=0}^{d}f_ix^i$ over $\mathbb{F}_q$ of least degree for which*

$$f\circ^q\alpha:=\sum_{i=0}^{d}f_i\alpha^{q^i}$$

is zero. We write $\mathrm{Ord}(\alpha)=f(x)$.

Notice that $(x^n-1)\circ^q\alpha=\alpha^{q^n}-\alpha=0$ for $\alpha\in\mathbb{F}_{q^n}$, hence the $\mathbb{F}_q-$Order of an element $\alpha\in\mathbb{F}_{q^n}$ is a polynomial of degree at most n. Also, it can be verified that such polynomial divides x^n-1. The following lemma gives a characterization of $k-$normal elements in $\mathbb{F}_{q^n}$:

Lemma 2.3 ([7], Theorem 3.2)**.** *Let $\alpha\in\mathbb{F}_{q^n}^*$. Then α is a $k-$normal element over $\mathbb{F}_q$ if and only if* $\deg(\mathrm{Ord}(\alpha))=n-k$.

Definition 2.4. *Let $f(x)$ be a monic polynomial with coefficients in $\mathbb{F}_q$.*
(a) The Euler Phi Function for polynomials over $\mathbb{F}_q$ is given by

$$\Phi_q(f)=\left|\left(\frac{\mathbb{F}_q[x]}{\langle f\rangle}\right)^*\right|,$$

where $\langle f\rangle$ is the ideal generated by $f(x)$ in $\mathbb{F}_q[x]$.

(b) The Mobius Function for polynomials over $\mathbb{F}_q$ is given by $\mu_q(f) = 0$ if $f(x)$ has some repeated irreducible factor over $\mathbb{F}_q$. Otherwise, $\mu_q(f) = (-1)^s$, where s is the number of irreducible factors of $f(x)$ over $\mathbb{F}_q$.

We have an interesting formula for the number of $k-$normal elements:

Lemma 2.5 ([7], Theorem 3.5)**.** *The number N_k of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is given by*

$$N_k = \sum_{\substack{h|x^n-1 \\ \deg(h)=n-k}} \Phi_q(h), \tag{1}$$

where the divisors are monic and polynomial division is over $\mathbb{F}_q$.

Since $\frac{x^n-1}{x-1}$ divides $x^n - 1$ and has degree $n - 1$, there are $1-$normal elements over any finite field extension. However, the sum in equality (1) can be empty: for instance, if $q = 7$ and $n = 5$, then

$$x^5 - 1 = (x - 1)(x^4 + x^3 + x^2 + x + 1)$$

is the factorization of $x^5 - 1$ into irreducible factors over $\mathbb{F}_7$; therefore there are no 2 or $3-$normal elements of $\mathbb{F}_{7^5}$ over $\mathbb{F}_7$. The existence of $k-$normal elements is not always ensured for generic values of k.

2.2. **Estimations.**

Here we present some bounds that will be useful in the next sections.

Lemma 2.6. *If $d(m)$ denotes the number of divisors of m, then for all $m \geq 3$,*

$$d(m) \leq m^{\frac{1.5379 \log 2}{\log\log m}} < m^{\frac{1.1}{\log\log m}}.$$

Proof. This inequality is a direct consequence of the result in [10]. □

Using the previous lemma we obtain the following estimation:

Proposition 2.7. *For any $m \geq 3$ and $x > 0$,*

$$\sum_{\substack{d|m \\ d\leq x}} \varphi(d) < x \cdot m^{\frac{1.1}{\log\log m}}.$$

Proof. Notice that, for all $m \geq 3$ and $x \geq 0$ we have

$$\sum_{\substack{d|m \\ d\leq x}} \varphi(d) \leq x \cdot \sum_{\substack{d|m \\ d\leq x}} 1 \leq x \cdot d(m),$$

and then the result follows from Lemma 2.6. □

Proposition 2.8. *For any positive integers k and $q \geq 2$ the following inequality holds:*

$$(q-1)^k \geq 4^{-k/q} q^k.$$

Proof. Notice that $a_n = (1-1/n)^n$ is an increasing sequence. Hence, for any $q \geq 2$ we have that $a_q \geq a_2 = 4^{-1}$ and then

$$(q-1)^k = ((q-1)^q)^{k/q} = (a_q \cdot q^q)^{k/q} \geq 4^{-k/q} q^k.$$

□

3. Existence questions for primitive $k-$normal elements

The Primitive Normal Basis Theorem states that for all pairs (q,n), where q is a power of a prime and n is a positive integer, there exists a normal basis $B = \{\alpha, \cdots, \alpha^{q^{n-1}}\}$ of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ such that α generates $\mathbb{F}_{q^n}^*$, i.e., α is primitive. A similar result was proved for $1-$normal elements:

Theorem 3.1 ([7], Theorem 5.10)**.** *Let $q = p^e$ be a prime power and n a positive integer not divisible by p. Assume that $n \geq 6$ if $q \geq 11$ and that $n \geq 3$ if $3 \leq q \leq 9$. Then there exists a primitive $1-$normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

We ask if this result can be generalized. For instance, under the condition that n is not divisible by p, can we guarantee the existence of primitive $k-$normal elements for some $k > 1$. The number $k-$normal elements can be zero for generic values of k, so any generalization of this result is under the hypothesis that $k-$normal elements actually exist.

In this section, as in [7], we apply the traditional method of Lenstra and Schoof in the characterization of primitive $k-$normal elements. First we need to recall the concept of *freeness*.

Definition 3.2. *(a) Assume that m divides $q^n - 1$. An element $\alpha \in \mathbb{F}_{q^n}$ is said to be $m-$free if for any d dividing m, the equality $\beta^d = \alpha$ with $\beta \in \mathbb{F}_{q^n}$ implies $d = 1$.*

(b) Assume that $M(x)$ is a monic divisor of $x^n - 1$. An element $\alpha \in \mathbb{F}_{q^n}$ is said to be $M(x)-$free if for any monic divisor $N(x)$ of $M(x)$, the equality $\alpha = N \circ^q \beta$ with $\beta \in \mathbb{F}_{q^n}$ implies $N(x) = 1$.

A simple calculation shows that $\alpha \in \mathbb{F}_{q^n}^*$ is primitive if and only if is $(q^n-1)-$free. Also, by definition, an element $\alpha \in \mathbb{F}_{q^n}$ is normal over $\mathbb{F}_q$ if and only if is $(x^n-1)-$free (in other words, α has $\mathbb{F}_q-$Order equal $x^n - 1$). In this context, the following lemma shows a relation between freeness and the $\mathbb{F}_q-$Order of an element:

Lemma 3.3 ([7], Theorem 5.4)**.** *Let α be any element in $\mathbb{F}_{q^n}$ and $f(x)$ its $\mathbb{F}_q-$Order. Then for any $g(x)$ dividing $x^n - 1$ the following are equivalent:*

(a) α is $g(x)-$free,

(b) $g(x)$ and $\frac{x^n-1}{f(x)}$ are coprime.

Under the condition that n is not divisible by the characteristic p of $\mathbb{F}_{q^n}$, in [[7] - Proposition 5.6], the authors give an interesting characterization of the elements in $\mathbb{F}_{q^n}$ whose $\mathbb{F}_q-$Order equals $\frac{x^n-1}{x-1}$. This characterization is one of the main ideas in the proof of Theorem 3.1. They notice that, in general, it is hard to characterize the elements of $\mathbb{F}_{q^n}$ with prescribed $\mathbb{F}_q-$Order. However, we can give a fair extension of that characterization:

Proposition 3.4. *Suppose that n is not divisible by the characteristic p of $\mathbb{F}_{q^n}$ and let $f_1(x), \cdots, f_s(x)$ be distinct monic irreducible factors of x^n-1 over $\mathbb{F}_q$. Then $\alpha \in \mathbb{F}_{q^n}$ has $\mathbb{F}_q-$Order equal*

$$T(x) := \frac{x^n-1}{f_1(x)\cdots f_s(x)},$$

if and only if α is $T(x)-$free but not $T_i(x)-$free for $1 \leq i \leq s$, where $T_i(x) = T(x)\cdot f_i(x)$.

Proof. Suppose that $\alpha \in \mathbb{F}_{q^n}$ has $\mathbb{F}_q-$Order equal $T(x)$. In particular, since n is not divisible by p, x^n-1 has no repeated irreducible factors. Therefore $\frac{x^n-1}{T(x)} = f_1(x)\cdots f_s(x)$ and $T(x)$ are coprime, while $f_1(x)\cdots f_s(x)$ and $T_i(x)$ are not coprime for $1 \leq i \leq s$. From Lemma 3.3, it follows that α is $T(x)-$free but not $T_i(x)-$free for any $1 \leq i \leq s$. Conversely, let $h(x) = \mathrm{Ord}(\alpha)$ and suppose that α is $T(x)-$free but not $T_i(x)-$free for $1 \leq i \leq s$. Again, from Lemma 3.3, it follows that $\frac{x^n-1}{h(x)}$ and $T(x)$ are coprime but $\frac{x^n-1}{h(x)}$ and $T_i(x)$ are not coprime. Using again that x^n-1 has no repeated irreducible factors, we conclude that $T(x)$ divides $h(x)$ but $T_i(x)$ does not divide $h(x)$. Since $x^n-1 = T(x)\cdot(f_1(x)\cdots f_s(x))$ and $h(x)$ divides x^n-1, it follows that $h(x) = T(x)$. □

3.1. **Existence of primitive $k-$normal elements.**

Here we establish an existence result on primitive $k-$normal elements. As in ([7] - Section 5.2), we use the *Lenstra-Schoof* method involving characteristic functions and character sums in order to characterize a particular class of primitive $k-$normal elements. Most of the techniques used here are similar to the ones present in [7]. For this reason we skip some details, which can be found in [5]. We adopt the notation of [5]:

Multiplicative Part: $\int\limits_{d|m} \eta_d$ represents the sum $\sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{(d)} \eta_d$, where μ and φ denote the Mobius and Euler functions for integers, respectively, η_d is a typical multiplicative character of $\mathbb{F}_{q^n}$ of order d, and the sum $\sum_{(d)} \eta_d$ runs through all the multiplicative characters of order d. If P denotes the set of $(q^n-1)-$free elements $\alpha \in \mathbb{F}_{q^n}$ (or, equivalently, of primitive elements) then the characteristic function $\mathbb{I}_P$ of P, satisfies:

$$\mathbb{I}_P(\omega) = \theta(q^n - 1) \int\limits_{d|q^n-1} \eta_d(\omega), \quad \omega \in \mathbb{F}_{q^n}, \tag{2}$$

where $\theta(q^n-1) = \frac{\varphi(q^n-1)}{q^n-1}$.

Additive Part: Let χ denote the canonical additive character on $\mathbb{F}_{q^n}$, i.e.

$$\chi(\omega) = \lambda\left(\sum_{i=0}^{n-1} \omega^{q^i}\right), \omega \in \mathbb{F}_{q^n},$$

where λ is the canonical additive character of $\mathbb{F}_q$ to $\mathbb{F}_p$.

If D is a monic polynomial dividing $x^n - 1$ over $\mathbb{F}_q$, a typical character χ_* of $\mathbb{F}_{q^n}$ of $\mathbb{F}_q-$Order D is one such that $\chi_*(D \circ^q \cdot)$ is the trivial additive character in $\mathbb{F}_{q^n}$ and D is minimal (in terms of degree) with this property. Let Δ_D be the set of all $\delta \in \mathbb{F}_{q^n}$ such that χ_δ has $\mathbb{F}_q-$Order D, where $\chi_\delta(\omega) = \chi(\delta\omega)$ for any $\omega \in \mathbb{F}_{q^n}$. For instance, $\Delta_1 = \{0\}$ and $\Delta_{x-1} = \mathbb{F}_q^*$.

Similarly, $\int\limits_{D|T} \chi_{\delta_D}$ represents the sum $\sum_{D|T} \frac{\mu_q(D)}{\Phi_q(D)} \sum_{(\delta_D)} \chi_{\delta_D}$, where μ_q and Φ_q denote the Mobius and Euler functions for polynomials over $\mathbb{F}_q$, respectively, χ_D denotes a typical additive character of $\mathbb{F}_{q^n}$ of $\mathbb{F}_q-$Order D and the sum $\sum_{(\delta_D)} \chi_{\delta_D}$ runs through all the additive characters whose $\mathbb{F}_q-$Order equals D, i.e., all $\delta_D \in \Delta_D$. If $g(x)$ is a monic polynomial dividing $x^n - 1$ and $S(g)$ denotes the set of $g(x)-$free elements of $\mathbb{F}_{q^n}$ then the characteristic function $\mathbb{I}_{S(g)}$ of $S(g)$, satisfies:

$$\mathbb{I}_{S(g)}(\omega) = \Theta(g) \int\limits_{D|g} \chi_{\delta_D}(\omega), \quad \omega \in \mathbb{F}_{q^n}, \tag{3}$$

where $\Theta(g) = \frac{\Phi_q(g)}{q^{\deg(g)}}$.

For the rest of this paper, we assume that q is a power of a prime p.

Lemma 3.5. *Suppose that n is not divisible by p and $T(x) := \frac{x^n-1}{f_1(x)\cdots f_s(x)}$, where $f_1(x), \cdots, f_s(x)$ are distinct monic irreducible factors of $x^n - 1$ over $\mathbb{F}_q$. Then I_T, the characteristic function for the set of elements in $\mathbb{F}_{q^n}$ whose $\mathbb{F}_q-$Order equals $T(x)$, satisfies the following:*

$$I_T(\omega) = \sum_{G|f_1\cdots f_s} \mu_q(G) \cdot \Theta(G \cdot T) \int\limits_{E|G\cdot T} \chi_{\delta_E}(\omega), \tag{4}$$

where G is monic and polynomial division is over $\mathbb{F}_q$.

Proof. From Proposition 3.4, the characteristic function for the set of elements of $\mathbb{F}_{q^n}$ with $\mathbb{F}_q-$Order equals $T(x)$ is described by the following

product:

$$I_T(\omega) = \mathbb{I}_{S(T)}(\omega) \prod_{i=1}^{s} (1 - \mathbb{I}_{S(T \cdot f_i(x))}(\omega)). \tag{5}$$

It follows directly from Lemma 3.3 that $\mathbb{I}_{S(A)}(\omega) \cdot \mathbb{I}_{S(B)}(\omega) = \mathbb{I}_{S(C)}(\omega)$ for any ω in $\mathbb{F}_{q^n}$, where A and B divide $x^n - 1$ and C is their least common multiple. Therefore, from equality (5), we obtain:

$$I_T(\omega) = \sum_{G | f_1 \cdots f_s} \mu_q(G) \cdot \mathbb{I}_{S(G \cdot T)}(\omega). \tag{6}$$

We combine equalities (6) and (3) and conclude the proof. □

f Denoting by $G_n(\eta, \chi)$ the Gauss sum $\sum_{w \in \mathbb{F}_{q^n}} \eta(\omega)\chi(\omega)$, we have the following identities:

(i) $G_n(1, 1) = q^n$,
(ii) $G_n(1, \chi) = G_n(\eta, 1) = 0$ if $\chi, \eta \neq 1$ and
(iii) $|G_n(\eta, \chi)| = q^{n/2}$ if $\chi, \eta \neq 1$,

where 1 denotes the trivial additive (or multiplicative) character. More details on characters and Gauss sums can be found in [9].

Proposition 3.6. *Suppose that n is not divisible by p and $T(x) := \frac{x^n - 1}{f_1(x) \cdots f_s(x)}$ has degree $n-k$, where $f_1(x), \cdots, f_s(x)$ are distinct monic irreducible factors of $x^n - 1$. Then the number N_T of primitive elements whose $\mathbb{F}_q-$Order equals $T(x)$, satisfies the following equality:*

$$\frac{N_T}{\theta(q^n - 1)\Theta(T)} = q^{n-k} + \sum_{G | f_1 \cdots f_s} \mu_q(G)\Theta(G) \int_{\substack{d | q^n - 1 \\ d \neq 1}} \int_{\substack{E | G \cdot T \\ E \neq 1}} G_n(\eta_d, \chi_{\delta_E}). \tag{7}$$

Proof. Combining the characteristic functions for the set of primitive elements and the elements with $\mathbb{F}_q-$Order equals $T(x)$ in (4) we obtain the following:

$$N_T = \sum_{w \in \mathbb{F}_{q^n}} \theta(q^n - 1) \left(\int_{d | q^n - 1} \eta_d(\omega) \right) \left(\sum_{G | f_1 \cdots f_s} \mu_q(G)\Theta(G \cdot T) \int_{E | G \cdot T} \chi_{\delta_E}(\omega) \right).$$

Since $\Theta(G \cdot T) = \Theta(G) \cdot \Theta(T)$, we obtain:

$$\frac{N_T}{\theta(q^n - 1)\Theta(T)} = \sum_{G | f_1 \cdots f_s} \mu_q(G)\Theta(G) \int_{d | q^n - 1} \int_{E | G \cdot T} G_n(\eta_d, \chi_{\delta_E}).$$

We look at the Gauss sums $G_n(\eta_d, \chi_{\delta_D})$: notice that $\eta_d \equiv 1$ only when $d = 1$, and $\chi_{\delta_E} \equiv 1$ only when $\delta_E = 0$, i.e., $E = 1$. Therefore, for each d dividing $q^n - 1$ and each E dividing $G \cdot T$, the following holds:

$$G_n(\eta_d, \chi_{\delta_E}) = \begin{cases} q^n & \text{for } d = E = 1, \\ 0 & \text{for } d = 1, E \neq 1 \text{ or } E = 1, d \neq 1, \\ \pm q^{n/2} & \text{otherwise.} \end{cases}$$

Using these identities and the fact that

$$\sum_{G|f_1\cdots f_s} \mu_q(G)\Theta(G) = \prod_{i=1}^{s}(1 - \Theta(f_i)) = \frac{1}{q^k},$$

we conclude the proof.

□

We are ready to state and prove our main result:

Theorem 3.7. *Let $n \geq 2$ be a positive integer not divisible by p. If k is a positive integer such that there exist $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ and*

$$k \leq \frac{n}{2}\left(1 - \frac{2.2}{\log\log(q^n - 1)} - \frac{\log 6}{\log q}\right),$$

then there exists a primitive $k-$normal element.

Proof. From Lemma 2.5, $x^n - 1$ is divisible by a polynomial $T(x)$ of degree $n-k$ over $\mathbb{F}_q$, hence $f(x) = \frac{x^n-1}{T(x)}$ has degree k. Let $f(x) = f_1(x)\cdots f_s(x)$ be the factorization of $f(x)$ into irreducible factors over $\mathbb{F}_q$. From Lemma 2.3, any element in $\mathbb{F}_{q^n}$ whose $\mathbb{F}_q-$Order equals $T(x)$ is $k-$*normal*. Therefore, if there is no primitive $k-$normal element in $\mathbb{F}_{q^n}$ then $N_T = 0$ in equation (7).

Let $W(t)$ be the number of square-free integers (or monic polynomials) dividing t. Since $x^n - 1$ has no repeated irreducible factors, $W(AB) = W(A)W(B)$ for any coprime polynomials A and B dividing $x^n - 1$ over $\mathbb{F}_q$. Applying estimates to equation (7) we obtain the following:

$$\begin{aligned} q^{n-k} &= \left|\frac{N_T}{\theta(q^n-1)\Theta(T)} - q^{n-k}\right| \\ &\leq W(q^n-1)\cdot \sum_{G|f_1\cdots f_s} q^{n/2}\cdot|\mu_q(G)\Theta(G)|W(G\cdot T) \\ &\leq q^{n/2}W(q^n-1)W(T)\cdot \sum_{G|f_1\cdots f_s} W(G). \end{aligned} \tag{8}$$

Notice that

$$\sum_{G|f_1\cdots f_s} W(G) = \prod_{i=1}^{s}(1 + W(f_i)) = 3^s$$

and $W(T) \leq 2^{\deg(T)} \leq 2^{n-s}$. Clearly $s \leq k$ and, from our hypothesis, $k < n/2$. Hence $s < n/2$ and then

$$W(T) \cdot \sum_{G|f_1 \cdots f_s} W(G) \leq 2^{n-s} \cdot 3^s < 6^{n/2} = q^{\frac{n}{2} \cdot \frac{\log 6}{\log q}}.$$

Since $q^n - 1 \geq 2^2 - 1 = 3$, from Lemma 2.6 we have

$$W(q^n - 1) \leq d(q^n - 1) < (q^n - 1)^{\frac{1.1}{\log\log(q^n-1)}} < q^{\frac{1.1n}{\log\log(q^n-1)}}.$$

Thus, under the assumption that

$$k \leq \frac{n}{2}\left(1 - \frac{2.2}{\log\log(q^n - 1)} - \frac{\log 6}{\log q}\right),$$

inequality (8) does not hold and we get a contradiction. □

Remark 3.8. *The function* $L(q,n) = \frac{n}{2}\left(1 - \frac{2.2}{\log\log(q^n-1)} - \frac{\log 6}{\log q}\right)$ *satisfies*

$$\lim_{q\to\infty} \frac{2L(q,n)}{n} = 1.$$

In particular, given $\varepsilon > 0$, *there exist a constant* M_ε *such that for* $q > M_\varepsilon$ *the following holds: if* $k \in \left[1, \frac{n}{2}(1-\varepsilon)\right]$ *and there exist* $k-$*normal elements of* $\mathbb{F}_{q^n}$ *over* $\mathbb{F}_q$ *then at least one of these* $k-$*normal elements is primitive.*

3.2. **A note on** $1-$**normal elements.**

In the case $k = 1$ and $T(x) = \frac{x^n-1}{x-1}$, after a simple reduction, equality (7) becomes:

$$\frac{N_T}{\theta(q^n-1)\Theta(T)} = q^{n-1} + \frac{1}{q} \int_{\substack{d|q^n-1 \\ d\neq 1}} \int_{\substack{E|T \\ E\neq 1}} G_n(\eta_d, \chi_{\delta_E}) - \left(1 - \frac{1}{q}\right) \int_{\substack{d|q^n-1 \\ d\neq 1}} \int_{E|T} G_n(\eta_d, \chi_{\delta_{(x-1)E}}).$$

If we suppose that $N_T = 0$, applying estimates to the previous equality in a similar way of (8), we obtain:

$$q^{n-1} \leq q^{n/2} W(T) W(q^n - 1) \cdot \left(\frac{1}{q} + \frac{q-1}{q}\right),$$

hence

$$q^{n/2-1} < W(T) W(q^n - 1).$$

The previous inequality is the final step in the proof of Theorem 3.1 in [7]. In this context, our work in this section generalizes that result on primitive $1-$normal elements.

4. Existence and order of $k-$normal elements

In the previous section we have proved an existence result for primitive $k-$normal elements and found out that our technique does not work for generic values of k. In this section we discuss a less restrictive question: the existence of $k-$normal elements with high multiplicative order in the group $\mathbb{F}_{q^n}^*$.

By high order elements we mean an element $\alpha \in \mathbb{F}_{q^n}^*$ such that $\text{ord}(\alpha)$, the multiplicative order of α in $\mathbb{F}_{q^n}^*$, is greater than any polynomial in n when n grows (see [6]). High order elements appears in the literature as an approach to primitive elements; they are useful in many applications such as the AKS primality test (see [1]). Most of the well known results on high order elements give exponential expressions in n as lower bounds: in *Artin-Schreier* extensions $K := \mathbb{F}_p[x]/(x^p - x - 1)$, the coset of x has multiplicative order at least $2^{2.54p}$ (see [11]).

The main idea here is the following: we find estimations for the number a_m of elements $\alpha \in \mathbb{F}_{q^n}$ with multiplicative order at most m and for the number N_k of $k-$normal elements. If $N_k \leq a_m$, we can guarantee the existence of $k-$normal elements with multiplicative order at least equal to m. We obtain the following:

Theorem 4.1. *Let q be a prime power and $\mathbb{F}_{q^n}$ be the finite field with q^n elements, where $n \geq 2$. Suppose that $k \leq n-1$ is a positive integer such that there exist $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. Then there exists some $k-$normal element whose multiplicative order in $\mathbb{F}_{q^n}^*$ is at least equal to*

$$\tau(k,q,n) := 4^{(k-n)/q} \cdot q^{n-k-\frac{1.1n}{\log\log(q^n-1)}}.$$

Proof. Notice that, if G is a finite cyclic group, for any d dividing $|G|$ there are exactly $\varphi(d)$ elements in G of order d. If $G = \mathbb{F}_{q^n}^*$ and A_m denotes the set of elements in $\mathbb{F}_{q^n}^*$ with multiplicative order at most m, since $q^n - 1 \geq 3$, from Proposition 2.7 we have the following inequality:

$$a_m := |A_m| = \sum_{\substack{d \mid q^n-1 \\ d \leq m}} \varphi(d) < m \cdot q^{\frac{1.1n}{\log\log(q^n-1)}}.$$

If B_k denotes the set of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$, by our assumption, $N_k := |B_k| \neq 0$. Hence, from Lemma 2.5, there exist some monic polynomial $h(x)$ of degree $n-k$ dividing $x^n - 1$ and $N_k \geq \Phi(h(x))$. If $h(x) = \prod_{i=1}^s f_i(x)$ is the factorization of $h(x)$ over $\mathbb{F}_q$ with $d_i = \deg f_i$, then $\sum_{i=1}^s d_i = n-k$ and

$$\Phi(h(x)) = \prod_{i=1}^{s}(q^{d_i} - 1) \geq \prod_{i=1}^{s}(q-1)^{d_i} = (q-1)^{n-k}.$$

Also, from Proposition 2.8 we have $(q-1)^{n-k} \geq 4^{(k-n)/q} q^{n-k}$, hence $N_k \geq 4^{(k-n)/q} q^{n-k}$. A simple calculation shows that if

$$m = \tau(k,q,n) = 4^{(k-n)/q} \cdot q^{n-k-\frac{1.1n}{\log\log(q^n-1)}},$$

then $a_m < N_k$ (this is the optimal value of m based on our estimations). Thus there exists some element in $B_k \cap (A_m)^c$ and we conclude the proof. □

Remark 4.2. *Notice that* $\tau(k,q,n) \geq q^{n(1-\varepsilon(q,n))-k}$, *where*

$$\varepsilon(q,n) = \frac{1.1}{\log\log(q^n-1)} + \frac{\log 4}{q\log q} \quad \textit{and} \quad \lim_{q\to\infty} \varepsilon(q,n) = 0.$$

In particular, for any $\varepsilon > 0$ *there exist a constant* M_ε *such that if* $q > M_\varepsilon$, *then* $\tau(k,q,n) > q^{n(1-\varepsilon)-k}$ *for every* $n \geq 2$ *and* $1 \leq k \leq n-1$.

As we have seen, most of the well known results on high order elements $\alpha \in \mathbb{F}_{q^n}$ give lower bounds like $\mathrm{ord}(\alpha) \geq c^n$, where $c > 1$ is a constant not depending on q (for more examples, see [3] and [4]). In general, those high-order elements are precisely exhibited, while the previous theorem concerns only on existence. On the other hand, under the condition that there exist $k-$normal elements, Theorem 4.1 proves that there is some of them having multiplicative order fairly greater than c^n when k is not close to n. For instance, given $0 < \gamma < 1/2$, for q sufficiently large we have $\tau(k,q,n) > q^{\gamma n}$ for all n and $k \in [1,(1-2\gamma)n]$.

4.1. **Existence of $k-$normal elements.**

We have discussed the existence of primitive and high-order $k-$normal elements in Theorems 3.7 and 4.1, under the condition that $k-$normal elements actually exist. As we have seen, the number of k-normal elements in finite field extensions can be zero for general values of k. In fact, Proposition 2.5 shows that the existence of $k-$normal elements is closely related to the factorization of $x^n - 1$ over $\mathbb{F}_q$. Here we give sufficient conditions on q and n to ensure the existence of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ for any $1 \leq k \leq n-1$. First, from the explicit factorization of $x^n - 1 \in \mathbb{F}_q[x]$ in [2], we have the following:

Lemma 4.3. *Suppose that* r *is a prime dividing* $q-1$. *Then for any positive integer* d *and any* $1 \leq k \leq r^d - 1$ *there exist a polynomial* $f(x)$ *of degree* k *dividing* $x^{r^d} - 1$ *over* $\mathbb{F}_q$ *such that* $f(1) \neq 0$.

Proof. Notice that r^d is not divisible by the characteristic of $\mathbb{F}_q$, hence $x^{r^d}-1$ has no repeated irreducible factors. The proof is divided in three cases:

(a) $r = 2$: in this case, $x^{2^d} - 1 = (x-1)\prod_{i=0}^{d-1}(x^{2^i}+1)$ (not necessarily the factorization into irreducible factors over $\mathbb{F}_q$). If $1 \leq k \leq 2^d - 1$, then $k = \sum_{i=0}^{d-1} s_i 2^i$ where $s_i \in \{0,1\}$. Therefore $f(x) = \prod_{i=0}^{d-1}(x^{2^i}+1)^{s_i}$

has degree k and divides $x^{2^d}-1$. Also, in this case, q is odd and then $f(1)=2^{\sum_{i=0}^{d-1}s_i}\neq 0$.

(b) $r>2$ and r^d divides $q-1$: in this case, $x^{r^d}-1=(x-1)\prod_{i=1}^{r^d-1}(x-\zeta_i)$ is the factorization of $x^{r^d}-1$ into irreducible factors over $\mathbb{F}_q$, where $\zeta_i\neq 1$ for all $1\leq i\leq r^d-1$. If $1\leq k\leq r^d-1$ then $f(x)=\prod_{i=1}^{k}(x-\zeta_i)$ has degree k, divides $x^{r^d}-1$ and satisfies $f(1)\neq 0$.

(c) $r>2$ and r^d does not divide $q-1$: in this case, 8 does not divide r^d and we use the result in (Corollary 3.2 (b), [2]); if r^s is the greatest power of r dividing $q-1$ then $1\leq s<d$ and the factorization of $x^{r^d}-1$ over $\mathbb{F}_q$ has r^s distinct irreducible factors of degree one and, for each $1\leq i\leq d-s$, $(r-1)r^{s-1}$ distinct irreducible factors of degree r^i. Let $1\leq k\leq r^d-1$ and $(a_{d-1}\cdots a_1a_0)_r$ be the $r-$adic representation of k with d digits. For each $s+1\leq i\leq d$, take $a_{i-1}r^{s-1}$ distinct irreducible factors of $x^{r^d}-1$ having degree r^{i-s} (this is possible since $0\leq a_i\leq r-1$ and $1\leq i-s\leq d-s$) and then take $D=\sum_{i=0}^{s-1}a_ir^i$ distinct linear factors of x^n-1, all different from $x-1$ (this is possible since $D\leq r^s-1$). The product of the chosen factors give us a polynomial $f(x)$ of degree $\sum_{i=0}^{d-1}a_ir^i=k$ such that $f(x)$ divides $x^{r^d}-1$ and satisfies $f(1)\neq 0$.

□

Using the previous lemma, we obtain the following existence result on $k-$normal elements:

Theorem 4.4. *Let $n\geq 2$ be a positive integer. Suppose that $rad(n)$ divides $p(q-1)$. Then for any $1\leq k\leq n-1$ there exist $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

Proof. From Proposition 2.5 it is enough to prove that for any $1\leq k\leq n-1$ there exists a polynomial of degree k dividing x^n-1. We will prove the last statement by induction on the number N of distinct prime factors of n. If $N=1$ and $n=p^t$ for some $t\geq 1$ the result is trivial, otherwise n is a power of a prime r dividing $q-1$ and then we use Lemma 4.3. Suppose that the statement is true for all positive integers up to $N\geq 1$ and that n has $N+1\geq 2$ distinct prime factors. We have two cases to consider:

(a) p divides n: we can write $n=p^un_0$, where $\gcd(p,n_0)=1$, n_0 has N distinct prime factors and $\mathrm{rad}(n_0)$ divides $q-1$. Let k be a positive integer between 1 and $n-1$. Then $k=an_0+b$, where $0\leq a<p^u$ and $0\leq b<n_0$. By the induction hypothesis, there exists a polynomial $g(x)$ of degree b dividing $x^{n_0}-1$ (if $b=0$, take $g(x)=1$). Therefore $f(x)=(x^{n_0}-1)^ag(x)$ has degree k and divides $(x^{n_0}-1)^{a+1}$, hence divides $(x^{n_0}-1)^{p^u}=x^n-1$.

(b) p does not divide n: this case is slight different from the previous one. We can write $n=r^un_1$ where r is a prime dividing $q-1$, $\gcd(r,n_1)=1$, n_1

has N distinct prime factors and $\mathrm{rad}(n_1)$ divides $q-1$. Let k be a positive integer between 1 and $n-1$ and write $k = an_1 + b$, where $0 \leq a < r^u$ and $0 \leq b < n_1$. From Lemma 4.3, there exists a polynomial $g(x)$ of degree a dividing $x^{r^u} - 1$ such that $g(1) \neq 0$ (if $a = 0$, take $g(x) = 1$). By the induction hypothesis, there exists a polynomial $h(x)$ of degree b dividing $x^{n_1} - 1$ (if $b = 0$, take $h(x) = 1$). Then $f(x) = g(x^{n_1})h(x)$ has degree k and both $h(x)$ and $g(x^{n_1})$ divide $x^n - 1$. To finish the proof, we show that $g(x^{n_1})$ and $h(x)$ are coprime: if there is some element α in the algebraic closure of $\mathbb{F}_q$ such that $g(\alpha^{n_1}) = h(\alpha) = 0$, since $h(x)$ divides $x^{n_1} - 1$, we have that $\alpha^{n_1} = 1$ and then $g(1) = 0$, a contradiction with the assumption $g(1) \neq 0$.

□

As a direct consequence of Theorems 3.7, 4.1 and 4.4 we have the following:

Corollary 4.5. *Suppose that q is a power of a prime p and $n \geq 2$ is a positive integer.*

(a) If $\mathrm{rad}(n)$ divides $q-1$, then for any $1 \leq k \leq \frac{n}{2}\left(1 - \frac{2.2}{\log\log(q^n-1)} - \frac{\log 6}{\log q}\right)$ there exist primitive $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.

(b) If $\mathrm{rad}(n)$ divides $p(q-1)$, then for any $1 \leq k \leq n-1$ there exists a $k-$normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ with multiplicative order in $\mathbb{F}_{q^n}^*$ at least

$$\tau(k,q,n) = 4^{(k-n)/q} \cdot q^{n-k-\frac{1.1n}{\log\log(q^n-1)}}.$$

In **Table 1** below, we give two numerical examples for the previous corollary. The first column gives the values of k, the second column gives an approximation of $\log \tau(k,q,n)$ (up to 2 decimal places) and the third one marks the values of k for which Theorem 3.7 ensures the existence of primitive $k-$normal elements.

Table 1: high-order and primitive $k-$normal elements.

$q = 2^{256}, n = 9$ $\log(q^n - 1) = 1,597.01$		
k	$\log \tau(k,q,n)$	pr.
1	1,181.40	*
2	1,003.95	*
3	826.50	*
4	649.06	
5	471.61	
6	294.17	
7	116.72	
8	-60.72	

$q = 3^{1001}, n = 8$ $\log(q^n - 1) = 8,797.69$		
k	$\log \tau(k,q,n)$	pr.
1	6,632.44	*
2	5,532.73	*
3	4,433.02	*
4	3,333.31	
5	2,233.60	
6	1,133.89	
7	34.18	

5. Conclusions and additional remarks

In this paper, we have partially answered two open problems on $k-$normal elements proposed in [7]. We prove existence results on high-order and primitive $k-$normal elements over finite fields. We have noticed that, in general, the number of $k-$normal elements can be zero and then we give sufficient conditions on q and n in order to guarantee the existence of $k-$normal elements for any $1 \leq k \leq n-1$.

We emphasize that the estimations presented in this paper are turning points in the proofs of Theorems 3.7 and 4.1 but they may not be sharp; we take these estimatons just because they give us satisfactory results and make the proofs more clean and simple. The upper bounds given in Lemma 2.6 and Proposition 2.7 are true for all positive integers $m \geq 3$. A more detailed study on the prime factorization of $m = q^n - 1$, combined with some tools in Analytic Number Theory, could yield slight improvements on our results.

References

[1] M. Agrawal, N. Kayal, N. Saxena, *Primes is in P*, Ann. of Math., 160 (2004) 781-793.

[2] F. E. Brochero Martínez, C. R. Giraldo Vergara, L. B. de Oliveira, *Explicit factorization of* $x^n - 1 \in \mathbb{F}_q[x]$, Des. Codes Cryptogr. 77 (2015) 277-286.

[3] F.E. Brochero Martínez, L. Reis, *Elements of high order in Artin-Schreier extensions of finite fields* $\mathbb{F}_q$, Finite Fields Appl. 41 (2016) 24-33.

[4] Q. Cheng, *Constructing finite field extensions with large order elements*, SIAM J. Discrete Math. 21 (2007), 726-730.

[5] S.D. Cohen, S. Huczynska. *The primitive normal basis theorem - without a computer*, Journal of the London Mathematical Society 67 (2003) 41-56.

[6] S. Gao, *Elements of provable high orders in finite fields*, Proc. Amer. Math. Soc. 127 (1999) 1615-1623.

[7] S. Huczynska, G.L. Mullen, D. Panario, and D. Thomson, *Existence and properties of* $k-$*normal elements over finite fields*, Finite Fields Appl. 24 (2013) 170-183.

[8] H.W. Lenstra, R. Schoof, *Primitive normal bases for finite fields*, Mathematics of Computation 48 (1987) 217-231.

[9] R. Lidl, H. Niederreiter, *Finite Fields: Encyclopedia of Mathematics and Its Applications*, vol. 20, 2nd ed. Cambridge University Pres, Cambridge, 1997.

[10] J.L. Nicolas, G. Robin, *Majorations explicites pour le nombre de diviseurs de N*, Canad. Math. Bull. 26 (1983) 485-492.

[11] J. F. Voloch, *On some subgroups of the multiplicative group of finite rings*, J. Théor. Nombres Bordeaux 16 (2004) 233-239.

Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, 30123-970, Brazil,

E-mail address: lucasreismat@gmail.com